

# Warming global temperatures may not affect carbon stored deep in northern peatlands, study says

December 13, 2016

👍 191

f

Like

6

G+

🐦

Tweet

📡

Submit

↑

submit

↓

reddit

★

Favorites

✉

Email

🖨

Print

📄

PDF



FSU Professor of Chemical Oceanography Jeff Chanton worked at a site operated by Oak Ridge National Laboratory to study the effects of global climate change. Credit: Jeff Chanton

Deep stores of carbon in northern peatlands may be safe from rising temperatures, according to a team of researchers from several U.S.-based institutions.

Ad closed by Google

And that is good news for now, the researchers said.

Florida State University research scientist Rachel Wilson and University of Oregon graduate student Anya Hopple are the first authors on a new study published today in *Nature Communications*. The study details experiments suggesting that carbon stored in peat—a highly organic material found in marsh or damp regions—may not succumb to the Earth's warming as easily as scientists thought.

That means if these northern peatlands—found in the upper half of the northern hemisphere—remain flooded, a substantial amount of carbon will not be released into the atmosphere.

"We do see some breakdown of peat on the surface, but not below 2 feet deep, where the bulk of the carbon is stored," Wilson said.

The study is part of a long-term look at how carbon stored in peat will respond to climate and environmental change. The team of researchers, led by Paul Hanson of the Oak Ridge National Laboratory, includes scientists from FSU, University of Oregon, Georgia Institute of Technology, the U.S. Department of Agriculture-Forest Service, Chapman University, Lawrence Livermore National Laboratory, Pacific Northwest National Laboratory and Oak Ridge National Laboratory.

Researchers ran four different temperature simulations—increasing the temperature of the peat by 2.25 degrees Celsius, 4.5 degrees Celsius, 6.25 degrees Celsius and 9 degrees Celsius—to see how it would respond to increased heat.

They found that the surface peat did emit more methane gas when warmed, but the deep peat did not break down and did not start emitting additional methane or carbon dioxide.

"If the release of greenhouse gases is not enhanced by temperature of the deep peat, that's great news because that means that if all other things remain as they are, the deep peat carbon remains in the soil," said Joel Kostka, professor of microbiology at Georgia Institute of Technology.

The Earth's soils contain 1,550 billion tons of organic carbon, and 500 billion tons of this carbon is stored in northern peatlands around the world. This quantity is roughly the same amount as carbon in the atmosphere.

Scientists have been anxious to learn how these northern peatlands will respond to warming because a tremendous amount of carbon could be released into the atmosphere.

Researchers worked at the Oak Ridge National Laboratory's experimental site known as SPRUCE in northern Minnesota to examine both surface peat and peat up to 6 feet deep. The majority of the carbon is stored deeper in the ground.

Large environmental chambers were constructed by the Oak Ridge team to enclose portions of the peatlands. Within these chambers, scientists simulated climate change effects such as higher temperatures and elevated carbon dioxide levels. They also took some of the deep peat back to their labs to heat in additional studies.

While scientists said they were surprised by the results, they also cautioned that this came only after one year of warming.

"There are the necessary caveats that this was only for one year, and the experiment is planned to run for a decade, and other ecosystem feedbacks may become important in greenhouse gas emissions," said Scott Bridgman, director of the Institute of Ecology and Evolution at University of Oregon and Hopple's adviser.

In the future, scientists also plan to look at how these peatlands respond to heightened carbon dioxide levels combined with the temperature increases.

"In the future, we'll be warmer, but we'll also have more carbon dioxide in the atmosphere, so we need to understand how these deep stores of peat, which have all this carbon, respond to these conditions," said Jeff Chanton, professor of oceanography at Florida State University.

🔍 Explore further: Peatland carbon storage is stabilized against catastrophic release of carbon

Journal reference: [Nature Communications](#)

30 shares

Provided by: [Florida State University](#)

🗉 feedback to editors




## MK Nanotechnology Products

Revisit Our Site Often For Updates On New Nanotechnologies And Products.

mknano.com



Related Stories



**Peatland carbon storage is stabilized against catastrophic release of carbon** 🕒 November 2, 2011  
Concerns that global warming may have a domino effect —unleashing 600 billion tons of carbon in vast expanses of peat in the Northern hemisphere and accelerating warming to disastrous proportions — may be less justified ...

**Smouldering peat fires may contribute to climate change** 🕒 September 24, 2013  
New research into smouldering wildfires in the UK has found that they could be a contributor to climate change.

**Boreal peatlands not a global warming time bomb** 🕒 June 10, 2015  
To some scientists studying climate change, boreal peatlands are considered a potential ticking time bomb. With huge stores of carbon in peat, the fear is that rising global temperatures could cause the release of massive ...

Recommended for you



**Researchers analyze 125 compounds from oil spilled in the Gulf of Mexico to determine their longevity** 🕒 December 19, 2016  
The oil discharged into the Gulf of Mexico following the explosion and sinking of the Deepwater Horizon (DWH) rig in 2010 contaminated more than 1,000 square miles of seafloor. The complexity of the event has made it difficult ...



**El Nino fueled Zika outbreak, new study suggests** 🕒 December 19, 2016  
Scientists at the University of Liverpool have shown that a change in weather patterns, brought on by the 'Godzilla' El Niño of 2015, fuelled the Zika outbreak in South America.



**The case of the missing diamonds** 🕒 December 19, 2016  
It all began innocently enough. Tyrone Daulton, a physicist with the Institute for Materials Science and Engineering at Washington University in St. Louis, was

User comments

Please sign in to add a comment. Registration is free, and takes less than a minute. [Read more](#)

email

password

Sign in

Click here to reset your password.

Sign in to get notified via email when new comments are made.

### More news stories



**Researchers analyze 125 compounds from oil spilled in the Gulf of Mexico to determine their longevity**  
The oil discharged into the Gulf of Mexico following the explosion and sinking of the Deepwater Horizon (DWH) rig in 2010 contaminated more than 1,000 square miles of seafloor. The complexity of the event has made it difficult ...




**El Nino fueled Zika outbreak, new study suggests**  
Scientists at the University of Liverpool have shown that a change in weather patterns, brought on by the 'Godzilla' El Niño of 2015, fuelled the Zika outbreak in South America.



**The case of the missing diamonds**  
It all began innocently enough. Tyrone Daulton, a physicist with the Institute for Materials Science and Engineering at Washington University in St. Louis, was



**Arctic lakes thawing earlier each year**  
Scientists from the University of Southampton have found Arctic lakes, covered with ice during the winter months, are melting earlier each spring.



**Satellites observe 'traffic jams' in Antarctic Ice Stream caused by tides**  
For the first time, researchers have closely observed how the ocean's tides can speed up or slow down the speed of glacial movement in Antarctica. The new data will help modelers better predict how glaciers will respond to ...



**Satellites help discover a jet stream in the Earth's core**  
A jet stream within the Earth's molten iron core has been discovered by scientists using the latest satellite data that helps create an 'x-ray' view of the planet.