

Thanks for the feedback! [Undo](#)
We'll use your feedback to review ads on this site.

https://www.sciencedaily.com/releases/2016/05/160530115540.htm



Science News

from research organizations

Print Email Share

Fish courtship pheromone uses the brain's smell pathway

Date: May 30, 2016
Source: RIKEN
Summary: Research has revealed that a molecule involved in fish reproduction activates the brain via the nose. The pheromone is released by female zebrafish and sensed by smell receptors in the noses of the males. The neural pathway and brain areas involved in transforming this molecular messenger into courtship behavior in fish were also identified by the researchers.

Share: Facebook Twitter Google+ LinkedIn + Email



RELATED TOPICS


- Mind & Brain
- > Neuroscience
 - > Perception
 - > Opium
- Plants & Animals
- > Fish
 - > Fisheries
 - > Mating and Breeding

RELATED TERMS

- > Sensory system
- > Oxytocin
- > Zebrafish
- > Itch
- > Excitotoxicity and cell damage
- > Bullying

Thanks for the feedback! [Undo](#)

We'll use your feedback to review ads on this site.



PGF2 α activates two ventromedial glomeruli in the zebrafish olfactory bulb, as revealed by neural activation marker (pERK; red). Green: GFP-expressing olfactory axons, blue: glomerular structure visualized with anti-SV2 antibody.

Credit: RIKEN

Research at the RIKEN Brain Science Institute in Japan has revealed that a molecule involved in fish reproduction activates the brain via the nose. The pheromone is released by female zebrafish and sensed by smell receptors in the noses of the males. The neural pathway and brain areas involved in transforming this molecular messenger into courtship behavior in fish were also identified and reported in *Nature Neuroscience* on May 30.

Prostaglandin F2 α (PGF2 α) is a reproductive hormone in female fish and mammals that is involved in ovulation and uterine contraction. In fish, it is also pheromone--a social and sexual signaling molecule used to attract the opposite sex--that is secreted by females in their urine. Males will normally swim toward even small concentrations of the molecule, but the researchers, led by Yoshihiro Yoshihara, found that male fish without a sense of smell were indifferent when PGF2 α was added to their tank.

PGF2 α synchronizes reproductive behaviors between female and male zebrafish, but how this happens was unknown. Having confirmed that sensory tissue responsible for smelling was needed for males to sense PGF2 α , researchers then found that the only type of neurons activated by it are ciliated olfactory sensory neurons. The team then searched for the receptor within these neurons that detects PGF2 α .

Contrary to expectations, the key players were not prostaglandin receptors. Molecular labeling revealed that PGF2 α only bound to two specific olfactory receptors. These olfactory receptors are evolutionarily quite different from prostaglandin receptors, and the same or corresponding olfactory receptor genes are present in other fish and mice, which indicates that a similar mechanism for reproductive communication might be present within other species.

The researchers also found that through these olfactory receptors, PGF2 α activates a direct, dedicated neural pathway to the areas of the brain that are responsible for eliciting courtship behavior in male fish. The ciliated olfactory sensory neurons send their signals to specific regions called glomeruli in the olfactory bulb of the brain, which in turn relay them to distinct forebrain areas. This "labeled line," in which circuits only transmit information about particular stimuli from a limited number of receptors, is also how the sense of taste functions. Hardwired pathways like this are common for innate behaviors, says Yoshihara, and it may have been an evolutionary accident that the PGF2 α molecule was well-matched to certain olfactory receptors, facilitating the use of the "smell pathway" for reproductive purposes.

Finally, the researchers tested the response to PGF2 α in male fish that were lacking the genes for one of the olfactory receptors they had identified. These fish weren't drawn to PGF2 α in their tank, spent less time chasing female fish, and were less successful at spawning. A smell receptor thus seems to be the gateway for PGF2 α into the male fish brain. Pheromone signaling works hand in hand with other senses like vision to bring about the courtship dance that increases a fish's chances of mating.

Story Source:

The above post is reprinted from materials provided by **RIKEN**. *Note: Materials may be edited for content and length.*

Journal Reference:

1. Yoichi Yabuki, Tetsuya Koide, Nobuhiko Miyasaka, Noriko Wakisaka, Miwa Masuda, Masamichi Ohkura, Junichi Nakai, Kyoshiro Tsuge, Soken Tsuchiya, Yukihiro Sugimoto, Yoshihiro Yoshihara. **Olfactory receptor for prostaglandin F2 α mediates male fish courtship behavior.** *Nature Neuroscience*, 2016; DOI: 10.1038/nn.4314

Cite This Page:

MLA APA Chicago

RIKEN. "Fish courtship pheromone uses the brain's smell pathway." ScienceDaily. ScienceDaily, 30 May 2016. <www.sciencedaily.com/releases/2016/05/160530115540.htm>.

Share This Page:

Facebook Twitter Google+ LinkedIn + Email

See Also

- | | |
|----------------------|------------------|
| 1 Pheromones Reviews | 4 Fish Posters |
| 2 Pheromone Spray | 5 Fish Furniture |
| 3 Primal Instinct | 6 Sea Fishing |

Recommended Articles

- It takes two to court: Pheromone that triggers mating found in mice
Stowers Institute for Medical Research, ScienceDaily, 2014
- The Brain Is Broadly Wired For Reproduction
Howard Hughes Medical Institute, ScienceDaily, 2005
- Tears for fears: Juvenile mice secrete a protective pheromone in their tears, blocking adult mating
Harvard Medical School, ScienceDaily, 2013
- Fruit Fly Pheromone Receptor First Ever Discovered Linked To Specific Sexual Behavior
Duke University Medical Center, ScienceDaily, 2003
- Researchers track neural circuits driving a fly's choice of a mate
Rockefeller University, ScienceDaily, 2015

Wake Up and Smell the iPhone [↗](#)
IEEE Pulse, 2015

Let There Be Light [↗](#)
IEEE Pulse, 2014

Forecast 2014 [↗](#)
IEEE Pulse, 2013

Scripps Florida study identifies memory suppressor gene that could hold key to new Alzheimer's disease treatments [↗](#)
MDLinx

Study offers insight on how a new class of antidepressants works [↗](#)
MDLinx

Related Stories

 Brain Circuit in Fruit Fly That Detects Anti-Aphrodisiac Uncovered


July 7, 2015 — New research has identified the neural circuit in the brain of the fruit fly (*Drosophila melanogaster*) that is responsible for detecting a taste pheromone, which controls the decision of male flies ... [read more »](#)

 Female Mice Are Able to Smell Male Pheromones Only When Ready to Mate

June 4, 2015 — An American study in mice reveals that hormones that dictate a female's attraction towards males do so in part by controlling her sense of smell. The findings provide an example of how hormones ... [read more »](#)

 It Takes Two to Court: Pheromone That Triggers Mating Found in Mice

July 25, 2014 — The functions of two classes of pheromone receptors have been identified by researchers, who have also found pheromones crucial to triggering the mating process in mice. "Interestingly, the ... [read more »](#)

 The Smoking Gun: Fish Brains and Nicotine

Dec. 9, 2013 — In researching neural pathways, it helps to establish an analogous relationship between a region of the human brain and the brains of more-easily studied animal species. New work hones in on one ... [read more »](#)

Thanks for the feedback! [Undo](#)

What was wrong with this ad?

- ☐ Inappropriate
- ☐ Repetitive
- ☐ Irrelevant




Strange & Offbeat

PLANTS & ANIMALS

 Dancing Hairs Alert Bees to Floral Electric Fields

 Hawk Moths Have Second Nose for Evaluating Flowers

Rapid Bone Growth and Underwater Breathing: Putting the Science of Harry Potter's Universe to the Test

 Tiny 'Vampires': Paleobiologist Finds Evidence of Predation in Ancient Microbial Ecosystems

EARTH & CLIMATE

 Methane-Producing Microbes in California Rocks

 In Changing Oceans, Cephalopods Are Booming

 Human-Eating Monster Crocodile May Be Florida's Newest Invasive Species

 Mighty Mealworms: Solution for Food Insecurity, Pollution

FOSSILS & RUINS

 New Horned Dinosaur Species With 'Spiked Shield'

 Ancient Irish Musical History Found in Modern India

 New Evidence Connects Dung Beetle Evolution to Dinosaurs

Medieval Women Better Dressed Than Men

Thanks for the feedback! [Undo](#)

We'll use your feedback to review ads on this site.



Print Email Share

Powered by Trend MD

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

-  Email Newsletters
-  RSS Feeds

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

-  Facebook
-  Twitter
-  Google+
-  LinkedIn

Mobile Apps

Get the latest news from ScienceDaily via our free mobile apps, available for download on the following platforms:

-  iPhone/iPad
-  Android

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

-  Leave Feedback
-  Contact Us