Sign up for WHO updates

Media centre
News
News releases
Previous years
Statements
Notes for the media
Events
Fact sheets
Features
Commentaries
Multimedia
Contacts

An estimated 12.6 million deaths each year are attributable to unhealthy environments

News release

Share Print

15 MARCH 2016 | GENEVA - An estimated 12.6 million people died as a result of living or working in an unhealthy environment in 2012 – nearly 1 in 4 of total global deaths, according to new estimates from WHO. Environmental risk factors, such as air, water and soil pollution, chemical exposures, climate change, and ultraviolet radiation, contribute to more than 100 diseases and injuries.

Noncommunicable diseases contribute to largest share of environment-related deaths

The second edition of the report, *"Preventing disease through healthy environments: a global assessment of the burden of disease from environmental risks"*, reveals that since the report was first published a decade ago, deaths due to noncommunicable diseases (NCDs), mostly attributable to air pollution (including exposure to second-hand tobacco smoke), amount to as much as 8.2 million of these deaths. NCDs, such as stroke, heart disease, cancers and chronic respiratory disease, now amount to nearly two-thirds of the total deaths caused by unhealthy environments.

At the same time, deaths from infectious diseases, such as diarrhoea and malaria, often related to poor water, sanitation and waste management, have declined. Increases in access to safe water and sanitation have been key contributors to this decline, alongside better access to immunization, insecticide-treated mosquito nets and essential medicines.

Healthier environment: healthier people

"A healthy environment underpins a healthy population," says Dr Margaret Chan, WHO Director-General. "If countries do not take actions to make environments where people live and work healthy, millions will continue to become ill and die too young."

The report emphasizes cost-effective measures that countries can take to reverse the upward trend of environment-related disease and deaths. These include reducing the use of solid fuels for cooking and increasing access to low-carbon energy technologies.

"There's an urgent need for investment in strategies to reduce environmental risks in our cities, homes and workplaces", said Dr Maria Neira, WHO Director, Department of Public Health, Environmental and Social Determinants of Health. "Such investments can significantly reduce the rising worldwide burden of cardiovascular and respiratory diseases, injuries, and cancers, and lead to immediate savings in healthcare costs."

Environmental risks take their greatest toll on young children and older people, the report finds, with children under 5 and adults aged 50 to 75 years most impacted. Yearly, the deaths of 1.7 million children under 5 and 4.9 million adults aged 50 to 75 could be prevented through better environmental management. Lower respiratory infections and diarrhoeal diseases mostly impact children under 5, while older people are most impacted by NCDs.

Burden of disease in WHO Regions

Regionally, the report finds, low- and middle-income countries in the WHO South-East Asia and Western Pacific Regions had the largest environment-related disease burden in 2012, with a total of 7.3 million deaths, most attributable to indoor and outdoor air pollution. Further regional statistics listed in the report include:

- 2.2 million deaths annually in African Region
- 847 000 deaths annually in Region of the Americas
- 854 000 deaths annually in Eastern Mediterranean Region
- 1.4 million deaths annually in European Region
- 3.8 million deaths annually in South-East Asia Region
- 3.5 million deaths annually in Western Pacific Region

Low- and middle-income countries bear the greatest environmental burden in all types of diseases and injuries, however for certain NCDs, such as cardiovascular diseases and cancers, the per capita disease burden can also be relatively high in high-income countries.

Top causes of environment-related deaths

Looking across more than 100 disease and injury categories, the report finds that the vast majority of environment-related deaths are due to cardiovascular diseases, such as stroke and ischaemic heart disease:

1. Stroke – 2.5 million deaths annually
2. Ischaemic heart disease – 2.3 million deaths annually
3. Unintentional injuries (such as road traffic deaths) – 1.7 million deaths annually
4. Cancers – 1.7 million deaths annually
5. Chronic respiratory diseases – 1.4 million deaths annually
6. Diarrhoeal diseases – 846 000 deaths annually
7. Respiratory infections – 567 000 deaths annually
8. Neonatal conditions – 270 000 deaths annually
9. Malaria – 259 000 deaths annually
10. Intentional injuries (such as suicides) – 246 000 deaths annually

Strategies to reduce environmental disease burden

"There's an urgent need for investment in strategies to reduce environmental risks in our cities, homes and workplaces", said Dr Maria Neira, WHO Director, Department of Public Health, Environmental and Social Determinants of Health. "Such investments can significantly reduce the rising worldwide burden of cardiovascular and respiratory diseases, injuries, and cancers, and lead to immediate savings in healthcare costs."

Environmental risks take their greatest toll on young children and older people, the report finds, with children under 5 and adults aged 50 to 75 years most impacted. Yearly, the deaths of 1.7 million children under 5 and 4.9 million adults aged 50 to 75 could be prevented through better environmental management. Lower respiratory infections and diarrhoeal diseases mostly impact children under 5, while older people are most impacted by NCDs.

Burden of disease in WHO Regions

Regionally, the report finds, low- and middle-income countries in the WHO South-East Asia and Western Pacific Regions had the largest environment-related disease burden in 2012, with a total of 7.3 million deaths, most attributable to indoor and outdoor air pollution. Further regional statistics listed in the report include:

- 2.2 million deaths annually in African Region
- 847 000 deaths annually in Region of the Americas
- 854 000 deaths annually in Eastern Mediterranean Region
- 1.4 million deaths annually in European Region
- 3.8 million deaths annually in South-East Asia Region
- 3.5 million deaths annually in Western Pacific Region

Low- and middle-income countries bear the greatest environmental burden in all types of diseases and injuries, however for certain NCDs, such as cardiovascular diseases and cancers, the per capita disease burden can also be relatively high in high-income countries.

Top causes of environment-related deaths

Looking across more than 100 disease and injury categories, the report finds that the vast majority of environment-related deaths are due to cardiovascular diseases, such as stroke and ischaemic heart disease:

1. Stroke – 2.5 million deaths annually
2. Ischaemic heart disease – 2.3 million deaths annually
3. Unintentional injuries (such as road traffic deaths) – 1.7 million deaths annually
4. Cancers – 1.7 million deaths annually
5. Chronic respiratory diseases – 1.4 million deaths annually
6. Diarrhoeal diseases – 846 000 deaths annually
7. Respiratory infections – 567 000 deaths annually
8. Neonatal conditions – 270 000 deaths annually
9. Malaria – 259 000 deaths annually
10. Intentional injuries (such as suicides) – 246 000 deaths annually

Strategies to reduce environmental disease burden

Publication

"Preventing disease through healthy environments: a global assessment of the burden of disease from environmental risks"

Fact file

10 facts on preventing disease through healthy environments

Commentary

Environments should improve not harm our health

Infographics



Arabic pdf, 1.64Mb
Chinese pdf, 1.68Mb
English pdf, 594kb
French pdf, 616kb
Russian pdf, 593kb
Spanish pdf, 1.78Mb

Related links

Public health, environmental and social determinants of health (PHE)
Water Safety Plans (WSPs)

Sign up for WHO updates

French pdf, 616kb
Russian pdf, 593kb
Spanish pdf, 1.78Mb

Related links

Public health, environmental and social determinants of health (PHE)
Water Safety Plans (WSPs)

Sign up for WHO updates

French pdf, 616kb
Russian pdf, 593kb
Spanish pdf, 1.78Mb

Related links

Public health, environmental and social determinants of health (PHE)
Water Safety Plans (WSPs)