

See fewer ads and support this site.

Contributor works on millions of sites and all of your devices.



Science News

from research organizations

Print Email Share

Desert mangroves are major source of carbon storage, study shows
Researchers estimate carbon storage capacity of coastal desert mangroves in Mexico

Date: March 28, 2016

Source: University of California - San Diego

Summary: Short, stunted mangroves living along the coastal desert of Baja California store up to five times more carbon below ground than their lush, tropical counterparts, researchers have found. The new study estimates that coastal desert mangroves, which only account for 1 percent of the land area, store nearly 30 percent of the region's belowground carbon.

Share: Facebook Twitter Google+ LinkedIn + Email

Thanks for the feedback! Undo

We'll use your feedback to review ads on this site.

FULL STORY



These are red mangroves (*Rhizophora mangle*) in a small forest patch in Baja California Peninsula.

Credit: Octavio Aburto / ILCP

Researchers found that short, stunted mangroves living along the coastal desert of Baja California store up to five times more carbon below ground than their lush, tropical counterparts. The new study led by researchers at Scripps Institution of Oceanography at UC San Diego estimates that coastal desert mangroves, which only account for one percent of the land area, store nearly 30 percent of the region's belowground carbon.

"Mangroves represent a thin layer between ocean and land, and yet we are seeing an ecosystem that is storing a lot of carbon in a very small area," said Paula Ezcurra, lead author of the study and a former researcher at Scripps.

The study, published in the *Proceedings of the National Academy of Sciences*, was the first to estimate carbon accumulated in the mangrove's sediment peat layer, and highlights the importance of better understanding the natural history conditions where mangroves live to more accurately estimate their global carbon storage capacity.

Mangroves are known to thrive in hot, salty, and muddy conditions along tropical coasts.

Scripps researchers collected sediment core samples from deep below the mangled roots at two desert lagoons?Bahia Magdalena and Balandra?and two lush, tropical lagoons?Marismas Nacionales and La Encrucijada? and calculated the carbon stored at the four locations.

The findings revealed a 2,000-year-old layer of non-decomposed roots, or peat, up to four meters (13 feet) under the desert mangroves. The peat acts like a sponge for stored atmospheric carbon, and provides a record of sea-level-rise history in the region.

The scientists discovered that desert mangroves restricted to rocky inlets on the rugged coast of Baja California have been growing over their own root remains over thousands of years to compensate for sea-level changes, accumulating a thick layer of peat below their roots. In contrast, mangroves in flat coastal floodplains have accumulated a thinner peat layer, since they have migrated over time as sea levels has risen, and increased river sedimentation has shifted the shoreline.

"The results show how these ecosystems have been adapting to climate change and sea-level rise for thousands of years," said Octavio Aburto-Oropeza, assistant professor at the Scripps Center for Marine Biodiversity and Conservation, who has been studying the ecosystem services of mangroves in Mexico, and in Baja California in particular, for more than 10 years. "Understanding the link between the natural history and carbon storage capacity of mangrove environments could give us insight into future climate change and sea-level rise."

Scientists estimate that mangrove forests are decreasing by three percent per year due to increased human activities along the shoreline in Mexico, and throughout the world.

Removal of mangroves result in the immediate release of thousands of years of stored carbon back into the atmosphere, and removes a natural ecosystem capable of sequestering carbon from human activity, thus causing further climate change impacts by returning a large amount of carbon into the atmosphere that has been stored underground for thousands of years.

"Our study helps to further quantify the value of mangrove forest ecosystems, and highlights the importance of mangrove conservation," said Exequiel Ezcurra, an adjunct professor at Scripps and professor of botany and plant sciences at University of California, Riverside. "Hopefully it will serve to underscore the urgent need for mangrove protection and for the serious appraisal of the environmental services provided by common-access resources."

Story Source:

The above post is reprinted from materials provided by **University of California - San Diego**. Note: Materials may be edited for content and length.

Journal Reference:

1. Octavio Aburto-Oropeza et al. **Coastal landforms and accumulation of mangrove peat increase carbon sequestration and storage.** *PNAS*, March 2016 DOI: 10.1073/pnas.1519774113

Cite This Page:

MLA APA Chicago

University of California - San Diego. "Desert mangroves are major source of carbon storage, study shows: Researchers estimate carbon storage capacity of coastal desert mangroves in Mexico." ScienceDaily. ScienceDaily, 28 March 2016. <www.sciencedaily.com/releases/2016/03/160328191832.htm>.

Thanks for the feedback! Undo

We'll use your feedback to review ads on this site.



Related Stories

Protected Areas Save Mangroves, Reduce Carbon Emissions

Sep. 14, 2015 — Protected areas not only keep significant swaths of Indonesia's shrinking mangrove habitats intact, but also prevent emissions of carbon dioxide that would have been released had these mangroves ... **read more »**



'Carbon Sink' Detected Underneath World's Deserts

July 28, 2015 — The world's deserts may be storing some of the climate-changing carbon dioxide emitted by human activities, a new study suggests. Massive aquifers underneath deserts could hold more carbon than ... **read more »**



When Mangroves No Longer Protect the Coastline

Feb. 1, 2013 — The mangrove forests in the Guyanas (French Guiana, Surinam and Guyana), which spread across the Orinoco and Amazon deltas, are among the most extensive in the world. This particular ecosystem, ... **read more »**

Mangroves Among the Most Carbon-Rich Forests in the Tropics; Coastal Trees Key to Lowering Greenhouse Gases

Apr. 4, 2011 — Coastal mangrove forests store more carbon than almost any other forest on Earth, according to a study conducted by a team of US Forest Service and university ... **read more »**

Amkov GearBest

Strange & Offbeat

PLANTS & ANIMALS



A World Map of Neanderthal and Denisovan Ancestry in Modern Humans



Creation of Minimal Cell With Just the Genes Needed for Independent Life



Moving Microswimmers Using Magnetic Vortices



Modified Maggots Could Help Human Wound Healing

EARTH & CLIMATE



City Birds Are Smarter Than Country Birds



Could Ireland's Ecosystems Cope If We Introduced St. Patrick's Scaly Foes?

First Successful Extraction of Ancient DNA from a Southern African Mummy



Unique Beak Evolved With Tool Use in New Caledonian Crow

FOSSILS & RUINS

Human Ancestors Explored 'out of Africa' Despite Impaired Nasal Faculties



More Ancient Viruses Lurk in Our DNA Than We Thought



The Linguistics of Signifying Time: The Human Gesture as Clock



Solving the Mystery of the Tully Monster

Introducing DE MONTFORT UNIVERSITY LEICESTER

With degrees in: Mechatronics Electrical & Electronic Engineering Computer Security

RANKED 54th IN THE UK

CLICK TO FIND OUT MORE AUSTON INSTITUTE OF MANAGEMENT

In Other News

from NewsDaily