

suki



strong safe solutions
for your skin problems
& sensitivities™

[learn more](#)

SIGN UP FOR FREE NEWSLETTER

EMAIL

[SIGN UP](#)

Join the Conversation
ENN
COMMUNITY

- original content
- post your comments

Stay updated with the ENN RSS feeds

Find us on: **facebook**

twitter

LinkedIn

ENN MEMBERS

- » [Our Editorial Affiliate Network](#)
- » [Partner Sites](#)

PRODUCTS & SERVICES

- » [Submit a News Release](#)
- » [Be a Part of ENN](#)

ADVERTISE WITH ENN

- » [Why Advertise with ENN](#)
- » [Contact ENN Sales Team](#)

ENN ARCHIVES

- » [News Archive](#)

ABOUT ENN

- » [History](#)
- » [Mission Statement](#)
- » [Our Staff](#)
- » [Editorial Philosophy](#)
- » [Contact ENN](#)
- » [ENN Terms](#)
- » [Fair Use Notice](#)

[Printer Friendly Version](#)

[ShareThis](#)

From: [Bristol University](#)
Published May 22, 2016 09:47 AM

Bristol University study shows how immune cells become activated

Immune cells play essential roles in the maintenance and repair of our bodies. When we injure ourselves, immune cells mount a rapid inflammatory response to protect us against infection and help heal the damaged tissue.

Lead researcher [Dr Helen Weavers](#), from the Faculty of Biomedical Sciences said: "While this immune response is beneficial for human health, many human diseases (including atherosclerosis, cancer and arthritis) are caused or aggravated by an overzealous immune response. A greater understanding of what activates the immune response is therefore crucial for the design of novel therapies to treat these inflammatory disorders.

"Our study found that immune cells must first become 'activated' by eating a dying neighbouring cell before they are able to respond to wounds or infection. In this way, immune cells build a molecular memory of this meal, which shapes their inflammatory behaviour."

The team's research, published in the journal *Cell*, used the fruit fly (*Drosophila melanogaster*) to study how a particular immune cell (the macrophage) becomes activated in order to respond to injury or infection. Using the fly allowed researchers to make time-lapse movies of the dynamic behaviour of the immune cells as they migrate within a living organism. It also allowed them to easily manipulate different genes and signalling pathways within the fly, to test which genes are important for immune cell behaviour.

Using genetics, the researchers dissected the mechanism by which the molecular memory is generated within the immune cell. Ingestion of the dying cell activates signalling via a calcium flash, which leads to an increase in the amount of an important damage receptor Draper in the immune cell. High levels of this receptor enable the 'primed' immune cell to sense the damage signals that entice them towards a wound during inflammation. Without this priming, the cells are blind to wounds and infections.


Image shows immune cells (green and red) migrating through the 3-D space (blue) within a living *Drosophila* embryo. Credit: Bristol University.

Read more at [Bristol University](#).

Like 0 [ShareThis](#)

RELATED ARTICLES

- » [Study points to new opportunities to fight cancer](#)
October 17, 2015 08:36 AM
- » [India stops further trials of HIV vaccine](#)
December 4, 2007 12:04 PM
- » [HIV Vaccine Target Could Solve Mutation Problem](#)
November 11, 2007 10:17 PM
- » [How the environment shapes our immune system](#)
January 15, 2015 04:32 PM



Get the ENN
mobile app
iOS Android

FEATURED
AFFILIATE ↗
ECOREPORT

View all the featured contributing content by this source

Lakes
Environmental
www.weblakes.com

Air Dispersion Models
AERMOD | CALPUFF | AUSTAL

Emissions Inventory
FETS | MOBILE | EMISSIONS

Risk Assessment
EcoRisk | IRAP-h

Meteorological Data
WRF | MM5 | Station

Training in Air Quality Modeling
"Real Expertise, and it shows!" -D.D. Madrid, Spain

