



Dust blowing through the Sahara Desert in 2012. Researchers have found that the plume was far less dusty between 5,000 and 11,000 years ago, containing only half the amount of dust that is transported today.

Image courtesy MODIS Rapid Response Team/Goddard Space Flight Center/NASA

Saharan dust in the wind

Scientists find huge reduction in African dust plume led to stronger Saharan monsoons 11,000 years ago.

Jennifer Chu | MIT News Office
November 23, 2016

Press Inquiries

RELATED

Paper: "Glacial to Holocene changes in trans-Atlantic Saharan dust transport and dust-climate feedbacks"

McGee Lab

Department of Earth, Atmospheric and Planetary Sciences

School of Science

ARCHIVES

Humans settled, set fire to Madagascar's forests 1,000 years ago

Intensity of desert storms may affect ocean phytoplankton

Stalagmites pinpoint drying of American West

A 'green' Sahara was far less dusty than today

Every year, trade winds over the Sahara Desert sweep up huge plumes of mineral dust, transporting hundreds of teragrams — enough to fill 10 million dump trucks — across North Africa and over the Atlantic Ocean. This dust can be blown for thousands of kilometers and The Sahara is the largest source of windblown dust to the Earth's atmosphere. But researchers from MIT, Yale University, and elsewhere now report that the African plume was far less dusty between 5,000 and 11,000 years ago, containing only half the amount of dust that is transported today.

In a paper published today in *Science Advances*, the researchers have reconstructed the African dust plume over the last 23,000 years and observed a dramatic reduction in dust beginning around 11,000 years ago. They say this weakened plume may have allowed more sunlight to reach the ocean, increasing its temperature by 0.15 degrees Celsius — a small but significant spike that likely helped whip up monsoons over North Africa, where climate at the time was far more temperate and hospitable than it is today.

McGee's co-authors include lead author Ross Williams, a former graduate student at MIT; along with Christopher Kinsley, Irit Tal, and David Ridley from MIT; Shineng Hu and Alexey Fedorov from Yale University; Richard Murray from Boston University; and Peter deMenocal from Columbia University.

A wet Sahara

Around 11,000 years ago, the Earth had just emerged from the last ice age and was beginning a new, interglacial epoch known as the Holocene. Geologists and archaeologists have found evidence that during this period the Sahara was much greener, wetter, and more livable than it is today.

"There was also extensive human settlement throughout the Sahara, with lifestyles that would never be possible today," McGee says. "Researchers at archaeological sites have found fish hooks and spears in the middle of the Sahara, in places that would be completely uninhabitable today. So there was clearly much more water and precipitation over the Sahara."

This evidence of wet conditions shows that the region experienced regular monsoon rains during the early Holocene. This was primarily due to the slow wobbling of Earth's axis, which exposed the Northern Hemisphere to more sunlight during summer; this, in turn, warmed the land and ocean and drew more water vapor — and precipitation — over North Africa. Increased vegetation in the Sahara may have also played a role, absorbing sunlight and heating the surface, drawing more moisture over the land.

"The mysterious thing is, if you try to simulate all these changes in these early and mid-Holocene climates, the models intensify the monsoons, but nowhere near the amounts suggested by the paleodata," McGee says. "One of the things not factored into these simulations is changes in windblown dust."

Tracking a dust plume

In their results published today, McGee and colleagues propose a reduction in African dust may indeed have contributed to increasing monsoon rains in the region. The researchers came to their conclusion after estimating the amount of long-range windblown dust emitted from Africa over the last 23,000 years, from the end of the last ice age to today.

They focused on dust transported long distances, as these particles are small and light enough to be lifted and carried through the atmosphere for days before settling thousands of kilometers away from their source. This fine-grained dust scatters incoming solar radiation, cooling the ocean's surface and potentially affecting precipitation patterns, depending on how much dust is in the air.

To estimate how the African dust plume has changed over thousands of years, the team looked for places where dust should accumulate rapidly. Dust can sink to the floor of open ocean, but there layers of sediment build up very slowly, at a rate of 1 centimeter every 1,000 years.

Places like the Bahamas, by contrast, accumulate sediment much more quickly, making it easier for scientists to determine the ages of particular sediment layers. What's more, it's been shown that most of the windblown dust that has accumulated in the Bahamas originated not from local regions such as the U.S., but from the Sahara.

Dust's climate role

McGee and his colleagues obtained sediment core samples from the Bahamas that were collected in the 1980s by scientists from the Woods Hole Oceanographic Institution. They brought the samples back to the lab and analyzed their chemical composition, including isotopes of thorium — an element that exists in windblown dust worldwide, at known concentrations.

They determined how much dust was in each sediment layer by measuring the primary isotope of thorium, and determined how fast it was accumulating by measuring the amount of a rare thorium isotope in each layer.

In this way, the team analyzed sediment layers from the last 23,000 years, and showed that around 16,000 years ago, toward the end of the last ice age, the dust plume was at its highest, lofting at least twice the amount of dust over the Atlantic, compared to today. However, between 5,000 and 11,000 years ago, this plume weakened significantly, with just half the amount of today's windblown dust.

Colleagues at Yale University then plugged their estimates into a climate model to see how such changes in the African dust plume would affect both ocean temperatures in the North Atlantic and overall climate in North Africa. The simulations showed that a drop in long-range windblown dust would raise sea surface temperatures by 0.15 degrees Celsius, drawing more water vapor over the Sahara, which would have helped to drive more intense monsoon rains in the region.

"The modeling showed that if dust had even relatively small impacts on sea surface temperatures, this could have pronounced impacts on precipitation and winds both in the north Atlantic and over North Africa," McGee says. Noting that the next key step is to reduce uncertainties in the modeling of dust's climate impacts, he adds: "We're not saying, the expansion of monsoon rains into the Sahara was caused solely by dust impacts. We're saying we need to figure out how big those dust impacts are, to understand both past and future climates."

Ina Tegen, a professor at the Leibniz Institute for Tropospheric Research in Germany, says the group's results suggest that "dust effects today may be considerable as well."

"Dust loads vary with changing climate, and due to the effects of dust on [solar] radiation, ice formation in clouds, and the carbon cycle, this may cause important climate feedbacks," says Tegen, who was not involved in the research. "The changing climate since the last ice age can be considered a 'natural laboratory' to study such effects. Understanding the past is the basis for predicting future changes with any confidence."

This research was supported, in part, by the National Science Foundation.

Topics:

Africa

Anthropology

Climate

Climate change

EAPS

Earth and atmospheric sciences

Environment

Geology

Research

School of Science

National Science Foundation (NSF)

MIT NEWS

November 29, 2016




3Q: Innovation districts

MIT professor explains how "productive neighborhoods" can remake cities.



Wires of Ice

MIT researchers discover astonishing behavior of water confined in carbon nanotubes.




Wi-Fi in Kendall Square

Free high-speed outdoor Internet service implemented by MIT and partners.



Marshall Scholars


Matthew Cavuto, Zachary Hulcher, Kevin Zhou, and Daniel Zuo will pursue two years of study in the U.K.



Prof. Thomas Kochan writes for The Huffington Post that a new social contract is needed in America to ensure that the economy works for everyone.

Kochan writes that "America needs to build a new social contract based on mutual respect and attuned to the needs of today's workforce and economy."

THE HUFFINGTON POST



Lauren Uhr: a brain researcher motivated by personal experience


MIT senior studies cognitive science and medicine from the vantage point of a person with dyslexia.



Enhancing education from pre-K to MIT and beyond




The Boston Globe



New method for analyzing crystal structure

Exotic materials called photonic crystals reveal their internal characteristics with new method.




Saharan dust in the wind

Scientists find huge reduction in African dust plume led to stronger Saharan monsoons 11,000 years ago.



The science of friction on graphene

Sliding on flexible graphene surfaces has been uncharted territory until now.



Nylon fibers made to flex like muscles

MIT engineers find a simple and inexpensive new approach to creating bending artificial muscle fibers.




Toward X-ray movies

Low-power tabletop source of ultrashort electron beams could replace car-size laboratory devices.




Biomarker could help guide cancer therapy, avoid drug resistance

Biologists link levels of Mana protein to breast cancer cells' sensitivity to chemotherapy.




At forum, MIT community tackles tough ethical questions of climate change

Why is it so hard for human beings to address climate change? What can motivate effective action?



Forging ahead on climate action



MIT Lincoln Laboratory garners six 2016 R&D 100 Awards

Brian Henderson: Experimenting with physics and policy

Creating new treatments for amblyopia

Brian Sullivan writes for Bloomberg about research affiliate Judah Cohen's 'Siberian Snow Theory,' which is based on the concept that the amount of snow covering the ground in northern Eurasia can be used to predict how cold winter will be in the northern hemisphere.

Sullivan writes that Cohen 'spies all the makings of an early, cold winter,' this year.

READ MORE

About This Website

This Website is maintained by the MIT News Office, part of the Office of Communications.

About the MIT News Office

MIT News Press Center

Press Inquiries

Filming Guidelines

Office of Communications

Contact Us

Terms of Use

RSS

Twitter

Facebook

Google+

Instagram

Flickr

YouTube

MIT Homepage

MIT Video

MIT Connect

MIT Resources

Events Calendar

About MIT

Admissions

Alumni

Education

Labs and Centers

Resources

Submit Campus News

Pitch a Story

Sign Up for Newsletter

Subscribe to Press Releases

Terms of Use