

facebook

Create a Facebook profile
Sign up for free and connect to the world

Sign up

https://www.sciencedaily.com/releases/2016/09/160907125002.htm

Science News

from research organizations

Print Email Share

Increased ocean acidification is due to human activities, say scientists

More anthropogenic carbon in the northeast Pacific means weaker shells for many marine species

Date: September 7, 2016

Source: Massachusetts Institute of Technology

Summary: Oceanographers report that the northeast Pacific Ocean has absorbed an increasing amount of anthropogenic carbon dioxide over the last decade, at a rate that mirrors the increase of carbon dioxide emissions pumped into the atmosphere.

Share:

RELATED TOPICS

Earth & Climate

- > Global Warming
- > Air Quality
- > Forest


Science & Society

- > Ocean Policy
- > Environmental Policies
- > Resource Shortage

Make your career in South Africa.


Enroll for BBA/MBA

Only a Few Seats Are Left



RELATED TERMS

- > Ocean acidification
- > Carbon dioxide
- > Carbon dioxide sink
- > Forest
- > Fossil fuel
- > Climate change mitigation



"The ocean has been the only true sink for anthropogenic emissions since the industrial revolution," says MIT graduate student Sophie Chu, pictured here. "Right now, it stores about 1/4 to 1/3 of the anthropogenic emissions from the atmosphere. We're expecting at some point the storage will slow down."

Credit: Zhaohui Aleck Wang/Woods Hole Oceanographic Institution

Oceanographers from MIT and Woods Hole Oceanographic Institution report that the northeast Pacific Ocean has absorbed an increasing amount of anthropogenic carbon dioxide over the last decade, at a rate that mirrors the increase of carbon dioxide emissions pumped into the atmosphere.

The scientists, led by graduate student Sophie Chu, in MIT's Department of Earth, Atmospheric, and Planetary Sciences, found that most of the anthropogenic carbon (carbon arising from human activity) in the northeast Pacific has lingered in the upper layers, changing the chemistry of the ocean as a result. In the past 10 years, the region's average pH has dropped by 0.002 pH units per year, leading to more acidic waters. The increased uptake in carbon dioxide has also decreased the availability of aragonite -- an essential mineral for many marine species' shells.

Overall, the researchers found that the northeast Pacific has a similar capacity to store carbon, compared to the rest of the Pacific. However, this carbon capacity is significantly lower than at similar latitudes in the Atlantic.

"The ocean has been the only true sink for anthropogenic emissions since the industrial revolution," Chu says. "Right now, it stores about 1/4 to 1/3 of the anthropogenic emissions from the atmosphere. We're expecting at some point the storage will slow down. When it does, more carbon dioxide will stay in the atmosphere, which means more warming. So it's really important that we continue to monitor this."

Chu and her colleagues have published their results in the *Journal of Geophysical Research: Oceans*.

Tippling the scales

The northeast Pacific, consisting of waters that flow from Alaska's Aleutian Islands to the tip of southern California, is considered somewhat of a climate canary -- sensitive to changes in ocean chemistry, and carbon dioxide in particular. The region sits at the end of the world's ocean circulation system, where it has collected some of the oldest waters on Earth and accumulated with them a large amount of dissolved inorganic carbon, which is naturally occurring carbon that has been respired by marine organisms over thousands of years.

"This puts the Pacific at this already heightened state of high carbon and low pH," Chu says.

Add enough atmospheric carbon dioxide into the mix, and the scales could tip toward an increasingly acidic ocean, which could have an effect first in sea snails called pteropods, which depend on aragonite (a form of calcium carbonate) to make their protective shells. More acidic waters can make carbonate less available to pteropods.

"These species are really sensitive to ocean acidification," Chu says. "It's harder for them to get enough carbonate to build their shells, and they end up with weaker shells, and have reduced growth rates."

Protecting shells

Chu and her colleagues originally set out to study the effects of ocean acidification on pteropods, rather than the ocean's capacity to store carbon. In 2012, the team embarked on a scientific cruise to the northeast Pacific, where they followed the same route as a similar cruise in 2001. During the month-long journey, the scientists collected samples of pteropods, as well as seawater, which they measured for temperature, salinity, and pH.

Upon their return, Chu realized that the data they collected could also be used to gauge changes in the ocean's anthropogenic carbon storage. Ordinarily, it's extremely difficult to tease out anthropogenic carbon in the ocean from carbon that naturally arises from breathing marine organisms. Both types of carbon are classified as dissolved inorganic carbon, and anthropogenic carbon in the ocean is miniscule compared to the vast amount of carbon that has accumulated naturally over millions of years.

To isolate anthropogenic carbon in the ocean and observe how it has changed through time, Chu used a modeling technique known as extended multiple linear regression -- a statistical method that models the relationships between given variables, based on observed data. The data she collected came from both the 2012 cruise and the previous 2001 cruise in the same region.

She ran a model for each year, plugging in water temperature, salinity, apparent oxygen utilization, and silicate. The models then estimated the natural variability in dissolved inorganic carbon for each year. That is, the models calculated the amount of carbon that should vary from 2001 to 2012, only based on natural processes such as organic respiration. Chu then subtracted the 2001 estimate from the 2012 estimate -- a difference that accounts for sources of carbon that are not naturally occurring, and are instead anthropogenic.

Sinking carbon

The researchers found that since 2001, the northeast Pacific has stored 11 micromoles per kilogram of anthropogenic carbon, which is comparable to the rate at which carbon dioxide has been emitted into the atmosphere. Most of this carbon is stored in surface waters. In the northern part of the region in particular, anthropogenic carbon tends to linger in shallower waters, within the upper 300 meters of the ocean. The southern region of the northeast Pacific stores carbon a bit deeper, within the top 600 meters.

Chu says this shallow storage is likely due to a subpolar gyre, or rotating current, that pushes water up from the deep, preventing surface waters from sinking. In contrast, others have observed that similar latitudes in the Atlantic have stored carbon much deeper, due to evaporation and mixing, leading to increased salinity and density, which causes carbon to sink.

The team calculated that the increase in anthropogenic carbon in the upper ocean caused a decrease in the region's average pH, making the ocean more acidic as a result. This acidification also had an effect on the region's aragonite, decreasing its saturation state over the last decade.

While the total amount of anthropogenic carbon appears to be increasing with each year, Chu says the rate at which the northeast Pacific has been storing carbon has remained relatively the same since 2001. That means that the region could still have a good amount of "room" to store carbon, at least for the foreseeable future. But already, her team and others are seeing in the acidification trends the ocean's negative response to the current rate of carbon storage.

"It would take hundreds of thousands of years for the ocean to absorb the majority of CO2 that humans have released into the atmosphere," Chu says. "But at the rate we're going, it's just way faster than anything can keep up with."

Story Source:

The above post is reprinted from materials provided by **Massachusetts Institute of Technology**. Note: Content may be edited for style and length.

Journal Reference:

- Sophie N. Chu, Zhaohui Aleck Wang, Scott C. Doney, Gareth L. Lawson, Katherine A. Hoering. **Changes in anthropogenic carbon storage in the Northeast Pacific in the last decade**. *Journal of Geophysical Research: Oceans*, 2016; 121 (7): 4618 DOI: 10.1002/2016JC011775

Cite This Page:

MLA APA Chicago

Massachusetts Institute of Technology. "Increased ocean acidification is due to human activities, say scientists: More anthropogenic carbon in the northeast Pacific means weaker shells for many marine species." ScienceDaily. ScienceDaily, 7 September 2016. <www.sciencedaily.com/releases/2016/09/160907125002.htm>.

facebook

Share your passion for sports, music, and more

Connect with Facebook

Sign Up

Related Stories

- Ocean Warming and Acidification Impact on Calcareous Phytoplankton and Reduce Its Ability to Sequester Atmospheric Carbon Dioxide
July 12, 2016 — Researchers warn of the negative impacts of rapid ocean warming and ocean acidification on coccolithophores, and consequently in the regulatory processes of atmospheric and ocean concentrations of ... [read more »](#)
- Climate Change Likely to Increase Black Carbon Input to the Arctic Ocean
Nov. 30, 2015 — The levels of black carbon in Arctic rivers has been determined by researchers who found that the input of black carbon to the Arctic Ocean is likely to increase with global ... [read more »](#)
- Carbon Dioxide-Spewing Volcano Drives Reef from Coral to Algae
Aug. 10, 2015 — A dramatic shift from vibrant coral communities to carpets of algae has been documented by researchers in remote Pacific Ocean waters where an underwater volcano spews carbon ... [read more »](#)

- Research Confirms How Global Warming Links to Carbon Emissions
Dec. 1, 2014 — Research has identified, for the first time, how global warming is related to the amount of carbon emitted. A team of researchers has derived the first theoretical equation to demonstrate that global ... [read more »](#)

Regenesys Business School
Awakening Potential

Make your career in South Africa. Enroll for BBA/MBA

Only a Few Seats Are Left

Strange & Offbeat

PLANTS & ANIMALS

- Trees Recognize Roe Deer by Saliva
- Researchers Debunk 'Five-Second Rule': Eating Food Off the Floor Isn't Safe
- Drones for Butterfly Conservation
- How Do Shark Teeth Bite? Reciprocating Saw, Glue Provide Answers

EARTH & CLIMATE

- Crab from the Chinese Pet Market Turns out to Be a New Species of a New Genus
- Tuning the Instrument: Spider Webs as Vibration Transmission Structures
- Fish 'Biowaste' Converted to Piezoelectric Energy Harvesters
- New Genus of Bacteria Found Living Inside Hydraulic Fracturing Wells

FOSSILS & RUINS

- Fossil Pollen 'Sneeze' Caught by Research Team
- A Mammoth Undertaking: Can De-Extinction Be Ecologically Responsible?
- 300 Teeth: Duck-Billed Dinosaurs Would Have Been Dentist's Dream
- Endangered Cuban Solenodon Evolved After the Extinction of Dinosaurs

Print Email Share

Powered by Google

CO2 Transmitters

cometsystem.com

World's rarest whale seen for the first time

sciencedaily.com

JYC Shafts Supplier

jyc.com.tw

Researchers Discover 'Hole' In Global...

sciencedaily.com

Heavy chocolate consumption may...

sciencedaily.com

Strongest material in world

graphene-uses.com

Why an ice age occurs every 100,000 years...

sciencedaily.com

Don't drink the (warm) water, study says

sciencedaily.com

Recommended Articles

- First evidence of ocean acidification affecting live marine creatures in the Southern Ocean
British Antarctic Survey, ScienceDaily, 2012
- Climate Change Seeps Into The Sea
NASA, ScienceDaily, 2008
- Global map of human-caused ocean acidification
The Earth Institute at Columbia University, ScienceDaily, 2014
- New research maps areas most vulnerable to ocean acidification
NOAA Headquarters, ScienceDaily, 2015
- New study reveals how sensitive US East Coast regions may be to ocean acidification
Woods Hole Oceanographic Institution, ScienceDaily, 2013

- Sequencing Study Reveals Pan Genome for Marine Phytoplankton Species
GenomeWeb, 2013
- Preservation of the Microbiological and Biochemical Quality of Raw Milk by Carbon Dioxide Addition: A Pilot-Scale Study
Ruas-Madiedo, Patricia et al., Journal of Food Protection
- Fate of Listeria monocytogenes in Shredded Cabbage Stored at 5 and 25°C under a Modified Atmosphere
Kallander, Kevin D. et al., Journal of Food Protection
- Microbiological and Chemical Changes of Spotted Shrimp (Pandalus platyceros) Stored Under Modified Atmospheres
Layrisse, Miguel E. et al., Journal of Food Protection
- Controlled-Atmosphere Storage of Pork Under Carbon Dioxide
Holley, R. A. et al., Journal of Food Protection

Powered by Trend MD

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

Email Newsletters

RSS Feeds

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

Facebook

Twitter

Google+

LinkedIn

Mobile Apps

Get the latest news from ScienceDaily via our free mobile apps, available for download on the following platforms:

iPhone/iPad

Android

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

Leave Feedback

Contact Us