



Centre for

Arctic Gas Hydrate, Environment and Climate



Search site ...



HOME

ABOUT

NEWS

RESEARCH

PUBLICATIONS

ARCTIC CHANGES

RESEARCH SCHOOL

INFRASTRUCTURES

CONTACT

24/03/2017

NORTHERN OCEANS PUMPED CO2 INTO THE ATMOSPHERE




Norwegian Sea acted as CO₂ source in the past. It pumped the greenhouse gas into the atmosphere instead of absorbing it, as it does today.

Tekst: Maja Sojtaric

At the same time the pH of the surface waters in these oceans decreased, making them more acidic. Both of these findings imply changes in ocean circulation and primary productivity as a result of natural climate changes of the time. The findings were recently published in *Nature Communications*.

Oceans changed function

Today the cold Arctic and Nordic Seas are especially efficient areas for uptake of CO₂ from the atmosphere. The oceans have been capable of mitigating some of the increase in greenhouse gas release resulting from human activities such as combustion of fossil fuels, by absorbing about 40% of the emitted CO₂




The illustration identifies the high-latitude North Atlantic as a significant CO₂ sink (The purple areas are the most efficient sinks, while red ones are sources of CO₂ in the modern ocean). The white star shows the location of the studied sediment core. The map was generated using data of Takahashi et al. Illustration: M. Ezat.

“Our research shows that areas in Norwegian Sea had changed their function on several occasions through the past 135 000 years: Instead of absorbing CO₂ from the air, they released more of the greenhouse gas into it.” says first author of the study Mohamed Ezat from Centre of Arctic Gas Hydrate, Environment and Climate (CAGE), Department of Geosciences at UiT The Arctic University of Norway.

First study of its kind from the Nordic Seas

Ice cores from Antarctica show that the amount of atmospheric CO₂ varied in pace with the shifting climate of ice ages and interglacial periods of the past.

“We always thought that oceans played a major role in these shifts, as it is the largest active CO₂ reservoir on this time scale. But it has remained unclear how and where in the ocean CO₂ was stored and released from”, says Ezat.




Micrefossils of single celled organisms record environmental changes of the time in their shells. Photo: Erik Thomsen and Tine Rasmussen.

Ezat and colleagues measured the boron isotopic composition of the fossil shells of near-surface dwelling, single celled organisms called foraminifera. These were collected from a marine sediment core, a record of the environment of the Norwegian sea spanning 135000 years. This period includes two warm interglacial periods, and one long lasting ice age characterised by abrupt climate changes.

“We saw that at the end of several of the severe cooling periods in the region, so-called Heinrich events, the ocean became more acidic and later released CO₂ into the atmosphere. These episodes of CO₂ pumping from the Nordic Seas coincide with times of increase in atmospheric CO₂,” says Ezat.

Measuring pH through thousands of years

“The variations in boron isotopes can tell us about the development in seawater pH through time and in turn give us information about the CO₂ concentration in the seawater.” explains co-author professor Tine L. Rasmussen, also from CAGE.



Mohamed Ezat in the laboratory of Lamont-Doherty Earth Observatory at Columbia University (USA), where he conducted the boron isotope measurements. Photo: Private

Doing so, the scientists were able to reconstruct the surface ocean pH and CO₂ in the Norwegian Sea in relation to past climate variations, when it was warmer or colder than today. Ezat and colleagues also tried to understand why the air-sea CO₂ exchange reversed in the Norwegian Sea during these times.

“We found that changes in primary productivity, input of terrestrial organic matter, and deep-water formation in the Nordic Seas, all contributed to the release of CO₂ from the ocean.” says Rasmussen

Never as acidic as today

The study shows that these seas had lower pH during the episodes of CO₂ release. This can however not be

compared to the extent of ocean acidification that we see happening today.

“Results of our study actually show that the sea surface pH throughout the last 135 000 years has never been as low as today in our study area. This is not an unexpected result. It is similar to previous studies conducted in other ocean areas. It does however add a body of evidence to the hypothesis that human activity is profoundly affecting the chemistry of our oceans,” Ezat says.

Scientists hope that the results will contribute to a better understanding of complex interactions between the ocean and atmosphere.

“In general, the more we learn about past changes in the Earth's climate system, the more accurate we hope we can predict the future.” says Ezat.

Reference: Ezat et al. Episodic release of CO₂ from the high-latitude North Atlantic Ocean during the last 135 kyr. *Nature Communications* 8, 2017. doi:10.1038/ncomms14498


EVENTS

International Training School - Bubbles 2017

06/06/2017 - 16/06/2017

Tromsø

Tweets by @CAGE_COE



Embed

View on Twitter

NEW PUBLICATIONS

Episodic release of CO₂ from the high-latitude North Atlantic Ocean during the last 135 kyr

Ecology of deep-sea benthic foraminifera in the North Atlantic during the last glaciation: food or temperature control

Ventilation history of Nordic Seas overflows during the last (de)glacial period revealed by species-specific benthic foraminiferal 14C dates

Large subglacial meltwater features in the central Barents Sea

The history and future trends of ocean warming-induced gas hydrate dissociation in the SW Barents Sea

Arctic megaslide at presumed rest



News Northern oceans pumped CO2 into the atmosphere

CAGE

Centre for Arctic Gas Hydrate, Environment and Climate
UiT The Arctic University of Norway
N-9019 Tromsø

Tel. +47 77 64 40 00

› All contact info

› All staff members



JOB OPENINGS

Please take a look at our career section to view vacant positions.

LINKS

- › The Research Council of Norway
- › UiT The Arctic University of Norway
- › Faculty of Science and Technology
- › Department of Geology
- › Arctic Portal
- › Database



GEOLOGICAL SURVEY OF NORWAY
NGU



The Research Council of Norway